

Global stability for coupled physics inverse problems. A case study

Giovanni Alessandrini



Università degli Studi di Trieste

Problèmes Inverses et Imagerie
12-13/02/2014
Institut Henri Poincaré

Introduction

Since the '80s, a dominant theme in Inverse Problems has been:

To image the interior of an object from measurements taken in its exterior

- overdetermined boundary data,
- scattering data.

With coupled physics IPs there is a shift to data associated to interior information.

Introduction

Since the '80s, a dominant theme in Inverse Problems has been:

To image the interior of an object from measurements taken in its exterior

- overdetermined boundary data,
- scattering data.

With coupled physics IPs there is a shift to data associated to interior information.

Introduction

Interior data may provide much better stability than inverse boundary problems, or inverse scattering problems.

Available results require nondegeneracy conditions on the solutions of the involved direct problems:

- Nonvanishing of solution.
- Nonvanishing of gradients.
- Nonvanishing of Jacobians.
- Nonvanishing of augmented Jacobians.

Introduction

Interior data may provide much better stability than inverse boundary problems, or inverse scattering problems.

Available results require nondegeneracy conditions on the solutions of the involved direct problems:

- Nonvanishing of solution.
- Nonvanishing of gradients.
- Nonvanishing of Jacobians.
- Nonvanishing of augmented Jacobians.

Introduction

Interior data may provide much better stability than inverse boundary problems, or inverse scattering problems.

Available results require nondegeneracy conditions on the solutions of the involved direct problems:

- Nonvanishing of solution.
- Nonvanishing of gradients.
- Nonvanishing of Jacobians.
- Nonvanishing of augmented Jacobians.

Introduction

Interior data may provide much better stability than inverse boundary problems, or inverse scattering problems.

Available results require nondegeneracy conditions on the solutions of the involved direct problems:

- Nonvanishing of solution.
- Nonvanishing of gradients.
- Nonvanishing of Jacobians.
- Nonvanishing of augmented Jacobians.

Introduction

Interior data may provide much better stability than inverse boundary problems, or inverse scattering problems.

Available results require nondegeneracy conditions on the solutions of the involved direct problems:

- Nonvanishing of solution.
- Nonvanishing of gradients.
- Nonvanishing of Jacobians.
- Nonvanishing of augmented Jacobians.

Global stability

Question. Is it possible to obtain global stability from measurements arising from arbitrary (nontrivial) solutions of the direct problem?

The model problem A problem arising in microwave imaging coupled with ultrasound, Triki (2010).

$$\Delta u + qu = 0 \text{ in } \Omega$$

Find $q \geq \text{constant} > 0$ given the local energy qu^2 and the boundary data $u|_{\partial\Omega}$.

Global stability

Question. Is it possible to obtain global stability from measurements arising from arbitrary (nontrivial) solutions of the direct problem?

The model problem A problem arising in microwave imaging coupled with ultrasound, Triki (2010).

$$\Delta u + qu = 0 \text{ in } \Omega$$

Find $q \geq \text{constant} > 0$ given the local energy qu^2 and the boundary data $u|_{\partial\Omega}$.

Global stability

Question. Is it possible to obtain global stability from measurements arising from arbitrary (nontrivial) solutions of the direct problem?

The model problem A problem arising in microwave imaging coupled with ultrasound, Triki (2010).

$$\Delta u + qu = 0 \text{ in } \Omega$$

Find $q \geq \text{constant} > 0$ given the local energy qu^2 and the boundary data $u|_{\partial\Omega}$.

Global stability

Question. Is it possible to obtain global stability from measurements arising from arbitrary (nontrivial) solutions of the direct problem?

The model problem A problem arising in microwave imaging coupled with ultrasound, Triki (2010).

$$\Delta u + qu = 0 \text{ in } \Omega$$

Find $q \geq \text{constant} > 0$ given the local energy qu^2 and the boundary data $u|_{\partial\Omega}$.

The full problem

Ammari, Capdeboscq, De Gournay, Rozanova-Pierrat, Triki (2011):

$$\operatorname{div}(a \nabla u) + k^2 q u = 0 \text{ in } \Omega$$

Find $a, q \geq \text{constant} > 0$ given the local energies $q u^2$, $a |\nabla u|^2$ (with several u 's and k 's!).

Here:

u = electric field, q = electric permittivity, a^{-1} = magnetic permeability.

The full problem

Ammari, Capdeboscq, De Gournay, Rozanova-Pierrat, Triki (2011):

$$\operatorname{div}(a \nabla u) + k^2 q u = 0 \text{ in } \Omega$$

Find $a, q \geq \text{constant} > 0$ given the local energies $q u^2$, $a |\nabla u|^2$ (with several u 's and k 's!).

Here:

u = electric field, q = electric permittivity, a^{-1} = magnetic permeability.

Introduction

An example

A priori
assumptions

Main Theorem

Stability for $|u|$

Quantitative
UCP

Concluding
remarks

End

Goals

- Stability of global type.
- Measurements for a single (nontrivial) solution u possibly sign changing.
- No spectral assumptions on $\Delta + q$.

Introduction

An example

A priori
assumptions

Main Theorem

Stability for $|u|$

Quantitative
UCP

Concluding
remarks

End

Goals

- **Stability of global type.**
- Measurements for a single (nontrivial) solution u possibly sign changing.
- No spectral assumptions on $\Delta + q$.

Goals

- Stability of global type.
- Measurements for a single (nontrivial) solution u possibly sign changing.
- No spectral assumptions on $\Delta + q$.

Goals

- Stability of global type.
- Measurements for a single (nontrivial) solution u possibly sign changing.
- No spectral assumptions on $\Delta + q$.

An example

In dimension $n = 1$, fix $0 < r < R$ and, for every $k = 1, 2, \dots$, set

$$q_k(x) = \begin{cases} A_k & \text{if } |x| < r, \\ 1 & \text{if } r \leq |x| \leq R, \end{cases}$$

where

$$A_k = \left(\frac{\pi}{2} + 2k\pi \right)^2 r^{-2}.$$

A solution to $u_{xx} + q_k u = 0$ in $(-R, R)$ is

$$u_k(x) = \begin{cases} \frac{1}{\sqrt{A_k}} \cos(\sqrt{A_k} x) & \text{if } |x| < r, \\ -\sin(|x| - r) & \text{if } r \leq |x| \leq R. \end{cases}$$

An example

In dimension $n = 1$, fix $0 < r < R$ and, for every $k = 1, 2, \dots$, set

$$q_k(x) = \begin{cases} A_k & \text{if } |x| < r, \\ 1 & \text{if } r \leq |x| \leq R, \end{cases}$$

where

$$A_k = \left(\frac{\pi}{2} + 2k\pi\right)^2 r^{-2}.$$

A solution to $u_{xx} + q_k u = 0$ in $(-R, R)$ is

$$u_k(x) = \begin{cases} \frac{1}{\sqrt{A_k}} \cos(\sqrt{A_k} x) & \text{if } |x| < r, \\ -\sin(|x| - r) & \text{if } r \leq |x| \leq R. \end{cases}$$

An example

we have

$$\|q_{2k} u_{2k}^2 - q_k u_k^2\|_\infty \leq 2$$

whereas, for any p , $1 \leq p \leq \infty$

$$\|q_{2k} - q_k\|_p \rightarrow \infty .$$

An example

we have

$$\|q_{2k} u_{2k}^2 - q_k u_k^2\|_\infty \leq 2$$

whereas, for any p , $1 \leq p \leq \infty$

$$\|q_{2k} - q_k\|_p \rightarrow \infty .$$

A priori assumptions

Given a bounded domain $\Omega \subset \mathbb{R}^n$ with Lipschitz boundary (in quantitative form!), we consider one solution $u \in W^{1,2}(\Omega) \cap C(\bar{\Omega})$ to

$$\Delta u + qu = 0 \text{ in } \Omega$$

where $q \in L^\infty(\Omega)$ is assumed to satisfy

$$0 < K^{-1} \leq q \leq K \text{ a.e. in } \Omega$$

for a given $K \geq 1$.

A priori assumptions

Energy bound. $E > 0$ is given such that:

$$\int_{\Omega} |\nabla u|^2 + u^2 \leq E^2 .$$

Nontriviality of the data. $H > 0$ is given such that:

$$\int_{\Omega} qu^2 \geq H^2 > 0 .$$

A priori data: K, E, H and Ω ($\text{diam}(\Omega)$, constants of its Lipschitz character).

Notation: For every $d > 0$: $\Omega_d = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > d\}$.

A priori assumptions

Energy bound. $E > 0$ is given such that:

$$\int_{\Omega} |\nabla u|^2 + u^2 \leq E^2 .$$

Nontriviality of the data. $H > 0$ is given such that:

$$\int_{\Omega} qu^2 \geq H^2 > 0 .$$

A priori data: K, E, H and Ω ($\text{diam}(\Omega)$, constants of its Lipschitz character).

Notation: For every $d > 0$: $\Omega_d = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > d\}$.

A priori assumptions

Energy bound. $E > 0$ is given such that:

$$\int_{\Omega} |\nabla u|^2 + u^2 \leq E^2 .$$

Nontriviality of the data. $H > 0$ is given such that:

$$\int_{\Omega} qu^2 \geq H^2 > 0 .$$

A priori data: K, E, H and Ω (**$\text{diam}(\Omega)$, constants of its Lipschitz character**).

Notation: For every $d > 0$: $\Omega_d = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > d\}$.

A priori assumptions

Energy bound. $E > 0$ is given such that:

$$\int_{\Omega} |\nabla u|^2 + u^2 \leq E^2 .$$

Nontriviality of the data. $H > 0$ is given such that:

$$\int_{\Omega} qu^2 \geq H^2 > 0 .$$

A priori data: K, E, H and Ω (**diam(Ω), constants of its Lipschitz character**).

Notation: For every $d > 0$: $\Omega_d = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > d\}$.

The main Theorem

Theorem. Let q_1, q_2 and the corresponding solutions u_1, u_2 satisfy the a priori assumptions and suppose that

$$\|q_1 u_1^2 - q_2 u_2^2\|_{L^\infty(\Omega)} \leq \varepsilon, \quad (1)$$

for a given $\varepsilon > 0$, and also

$$\||u_1| - |u_2|\|_{L^\infty(\partial\Omega)} \leq \sqrt{K\varepsilon}. \quad (2)$$

Then, for every $d > 0$, there exists $\eta \in (0, 1)$ and $C > 0$, only depending on d and on the a priori data such that

$$\|q_1 - q_2\|_{L^2(\Omega_d)} \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^\eta,$$

Note: If we know $q_1 = q_2$ near $\partial\Omega$, then (1) \Rightarrow (2).

The main Theorem

Theorem. Let q_1, q_2 and the corresponding solutions u_1, u_2 satisfy the a priori assumptions and suppose that

$$\|q_1 u_1^2 - q_2 u_2^2\|_{L^\infty(\Omega)} \leq \varepsilon, \quad (1)$$

for a given $\varepsilon > 0$, and also

$$\| |u_1| - |u_2| \|_{L^\infty(\partial\Omega)} \leq \sqrt{K\varepsilon}. \quad (2)$$

Then, for every $d > 0$, there exists $\eta \in (0, 1)$ and $C > 0$, only depending on d and on the a priori data such that

$$\|q_1 - q_2\|_{L^2(\Omega_d)} \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^\eta,$$

Note: If we know $q_1 = q_2$ near $\partial\Omega$, then (1) \Rightarrow (2).

The main Theorem

Theorem. Let q_1, q_2 and the corresponding solutions u_1, u_2 satisfy the a priori assumptions and suppose that

$$\|q_1 u_1^2 - q_2 u_2^2\|_{L^\infty(\Omega)} \leq \varepsilon, \quad (1)$$

for a given $\varepsilon > 0$, and also

$$\||u_1| - |u_2|\|_{L^\infty(\partial\Omega)} \leq \sqrt{K\varepsilon}. \quad (2)$$

Then, for every $d > 0$, there exists $\eta \in (0, 1)$ and $C > 0$, only depending on d and on **the a priori data** such that

$$\|q_1 - q_2\|_{L^2(\Omega_d)} \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^\eta,$$

Note: If we know $q_1 = q_2$ near $\partial\Omega$, then (1) \Rightarrow (2).

The main Theorem

Theorem. Let q_1, q_2 and the corresponding solutions u_1, u_2 satisfy the a priori assumptions and suppose that

$$\|q_1 u_1^2 - q_2 u_2^2\|_{L^\infty(\Omega)} \leq \varepsilon, \quad (1)$$

for a given $\varepsilon > 0$, and also

$$\||u_1| - |u_2|\|_{L^\infty(\partial\Omega)} \leq \sqrt{K\varepsilon}. \quad (2)$$

Then, for every $d > 0$, there exists $\eta \in (0, 1)$ and $C > 0$, only depending on d and on **the a priori data** such that

$$\|q_1 - q_2\|_{L^2(\Omega_d)} \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^\eta,$$

Note: If we know $q_1 = q_2$ near $\partial\Omega$, then (1) \Rightarrow (2).

Main tools

Theorem A (Stability for $|u|$)

There exists $C > 0$, only depending on K, E and Ω , such that

$$\int_{\Omega} ||u_1| - |u_2||^3 \leq C\varepsilon .$$

Theorem B (Integrability of $|u|^{-\delta}$) For every $d > 0$, there exists $\rho > 1, C > 0$, only depending on K, E, H and Ω , such that

$$\int_{\Omega_d} |u_1|^{-\frac{2}{\rho-1}} \leq C .$$

Note: This is a form of quantitative estimate of unique continuation.

Main tools

Theorem A (Stability for $|u|$)

There exists $C > 0$, only depending on K, E and Ω , such that

$$\int_{\Omega} ||u_1| - |u_2||^3 \leq C\varepsilon .$$

Theorem B (Integrability of $|u|^{-\delta}$) For every $d > 0$, there exists $\rho > 1, C > 0$, only depending on K, E, H and Ω , such that

$$\int_{\Omega_d} |u_1|^{-\frac{2}{\rho-1}} \leq C .$$

Note: This is a form of quantitative estimate of unique continuation.

Main tools

Theorem A (Stability for $|u|$)

There exists $C > 0$, only depending on K, E and Ω , such that

$$\int_{\Omega} ||u_1| - |u_2||^3 \leq C\varepsilon .$$

Theorem B (Integrability of $|u|^{-\delta}$) For every $d > 0$, there exists $\rho > 1, C > 0$, only depending on K, E, H and Ω , such that

$$\int_{\Omega_d} |u_1|^{-\frac{2}{\rho-1}} \leq C .$$

Note: This is a form of **quantitative estimate of unique continuation**.

Proof of the main theorem

$$\begin{aligned}(q_1 - q_2)u_1^2 &= q_2(u_2^2 - u_1^2) + (q_1u_1^2 - q_2u_2^2) = \\ &= q_2(|u_2| + |u_1|)(|u_2| - |u_1|) + (q_1u_1^2 - q_2u_2^2)\end{aligned}$$

hence

$$\int_{\Omega} |q_1 - q_2|u_1^2 \leq K \| |u_1| + |u_2| \|_{L^{3/2}(\Omega)} \| |u_1| - |u_2| \|_{L^3(\Omega)} + |\Omega|\varepsilon$$

and, by Theorem A,

$$\int_{\Omega} |q_1 - q_2|u_1^2 \leq C(\varepsilon^{1/3} + \varepsilon).$$

Proof of the main theorem

$$\begin{aligned}(q_1 - q_2)u_1^2 &= q_2(u_2^2 - u_1^2) + (q_1u_1^2 - q_2u_2^2) = \\ &= q_2(|u_2| + |u_1|)(|u_2| - |u_1|) + (q_1u_1^2 - q_2u_2^2)\end{aligned}$$

hence

$$\int_{\Omega} |q_1 - q_2|u_1^2 \leq K \| |u_1| + |u_2| \|_{L^{3/2}(\Omega)} \| |u_1| - |u_2| \|_{L^3(\Omega)} + |\Omega|\varepsilon$$

and, by Theorem A,

$$\int_{\Omega} |q_1 - q_2|u_1^2 \leq C(\varepsilon^{1/3} + \varepsilon).$$

Proof of the main theorem

Now, by Hölder's inequality, for any $\delta > 0$

$$\int_{\Omega_d} |q_1 - q_2|^{\frac{\delta}{\delta+2}} \leq \left(\int_{\Omega_d} |u_1|^{-\delta} \right)^{\frac{2}{\delta+2}} \left(\int_{\Omega} |q_1 - q_2| u_1^2 \right)^{\frac{\delta}{\delta+2}}$$

and choosing $\delta = \frac{2}{\rho-1}$, by Theorem B

$$\int_{\Omega_d} |q_1 - q_2|^{\frac{\delta}{\delta+2}} \leq C \left(\int_{\Omega} |q_1 - q_2| u_1^2 \right)^{\frac{\delta}{\delta+2}}.$$

Recalling $K^{-1} \leq q_i \leq K$ we arrive at

$$\int_{\Omega_d} |q_1 - q_2|^2 \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^{\frac{\delta}{\delta+2}}.$$

Proof of the main theorem

Now, by Hölder's inequality, for any $\delta > 0$

$$\int_{\Omega_d} |q_1 - q_2|^{\frac{\delta}{\delta+2}} \leq \left(\int_{\Omega_d} |u_1|^{-\delta} \right)^{\frac{2}{\delta+2}} \left(\int_{\Omega} |q_1 - q_2| u_1^2 \right)^{\frac{\delta}{\delta+2}}$$

and choosing $\delta = \frac{2}{p-1}$, by Theorem B

$$\int_{\Omega_d} |q_1 - q_2|^{\frac{\delta}{\delta+2}} \leq C \left(\int_{\Omega} |q_1 - q_2| u_1^2 \right)^{\frac{\delta}{\delta+2}}.$$

Recalling $K^{-1} \leq q_i \leq K$ we arrive at

$$\int_{\Omega_d} |q_1 - q_2|^2 \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^{\frac{\delta}{\delta+2}}.$$

Proof of the main theorem

Now, by Hölder's inequality, for any $\delta > 0$

$$\int_{\Omega_d} |q_1 - q_2|^{\frac{\delta}{\delta+2}} \leq \left(\int_{\Omega_d} |u_1|^{-\delta} \right)^{\frac{2}{\delta+2}} \left(\int_{\Omega} |q_1 - q_2| u_1^2 \right)^{\frac{\delta}{\delta+2}}$$

and choosing $\delta = \frac{2}{p-1}$, by Theorem B

$$\int_{\Omega_d} |q_1 - q_2|^{\frac{\delta}{\delta+2}} \leq C \left(\int_{\Omega} |q_1 - q_2| u_1^2 \right)^{\frac{\delta}{\delta+2}}.$$

Recalling $K^{-1} \leq q_i \leq K$ we arrive at

$$\int_{\Omega_d} |q_1 - q_2|^2 \leq C \left(\varepsilon^{1/3} + \varepsilon \right)^{\frac{\delta}{\delta+2}}.$$

Stability for $|u|$

Denote $N_i = \{u_i = 0\}$, $i = 1, 2$. Let Ω_j , $j = 1, 2, \dots$ be the connected components of $\Omega \setminus (N_1 \cup N_2)$.

For each j we may split

$$\partial\Omega_j = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2,$$

where

$$\Gamma_0 = \partial\Omega_j \cap \partial\Omega, \Gamma_1 = \partial\Omega_j \cap N_1, \Gamma_2 = \partial\Omega_j \cap N_2.$$

By assumption, on Γ_0 we have $\||u_1| - |u_2|\| \leq \sqrt{K\varepsilon}$, while, on Γ_1 , $q_2 u_2^2 \leq \varepsilon$ and on Γ_2 , $q_1 u_1^2 \leq \varepsilon$. Hence, on $\partial\Omega_j$ we have

$$\||u_1| - |u_2|\| \leq \sqrt{K\varepsilon},$$

Stability for $|u|$

Denote $N_i = \{u_i = 0\}$, $i = 1, 2$. Let Ω_j , $j = 1, 2, \dots$ be the connected components of $\Omega \setminus (N_1 \cup N_2)$.

For each j we may split

$$\partial\Omega_j = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2,$$

where

$$\Gamma_0 = \partial\Omega_j \cap \partial\Omega, \Gamma_1 = \partial\Omega_j \cap N_1, \Gamma_2 = \partial\Omega_j \cap N_2.$$

By assumption, on Γ_0 we have $\||u_1| - |u_2|\| \leq \sqrt{K\varepsilon}$, while, on Γ_1 , $q_2 u_2^2 \leq \varepsilon$ and on Γ_2 , $q_1 u_1^2 \leq \varepsilon$. Hence, on $\partial\Omega_j$ we have

$$\||u_1| - |u_2|\| \leq \sqrt{K\varepsilon},$$

Stability for $|u|$

Denote $N_i = \{u_i = 0\}$, $i = 1, 2$. Let Ω_j , $j = 1, 2, \dots$ be the connected components of $\Omega \setminus (N_1 \cup N_2)$.

For each j we may split

$$\partial\Omega_j = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2,$$

where

$$\Gamma_0 = \partial\Omega_j \cap \partial\Omega, \Gamma_1 = \partial\Omega_j \cap N_1, \Gamma_2 = \partial\Omega_j \cap N_2.$$

By assumption, on Γ_0 we have $\||u_1| - |u_2|\| \leq \sqrt{K\varepsilon}$, while, on Γ_1 , $q_2 u_2^2 \leq \varepsilon$ and on Γ_2 , $q_1 u_1^2 \leq \varepsilon$. Hence, on $\partial\Omega_j$ we have

$$\||u_1| - |u_2|\| \leq \sqrt{K\varepsilon},$$

Stability for $|u|$

W.l.o.g. we may assume $u_1, u_2 > 0$ in Ω_j . Set

$$\varphi^+ = [u_1 - u_2 - 2\sqrt{K\varepsilon}]^+, \varphi^- = [u_2 - u_1 - 2\sqrt{K\varepsilon}]^+.$$

Note that $\varphi^\pm \in W_0^{1,2}(\Omega_j) \cap C(\overline{\Omega_j})$ and use $\psi_j^\pm = \varphi^\pm u_j$ as test functions in the weak formulation of $\Delta u_j + q_j u_j = 0$. We arrive at

$$\int_{\Omega_j} (|u_1| + |u_2|)(|u_1| - |u_2|)^2 \leq C \int_{\Omega_j} (|u_1| + |u_2|)\varepsilon,$$

Adding up w.r.t. j , and using the energy bound,

$$\int_{\Omega} ||u_1| - |u_2||^3 \leq \int_{\Omega} (|u_1| + |u_2|)(|u_1| - |u_2|)^2 \leq C\varepsilon,$$

Stability for $|u|$

W.l.o.g. we may assume $u_1, u_2 > 0$ in Ω_j . Set

$$\varphi^+ = [u_1 - u_2 - 2\sqrt{K\varepsilon}]^+, \varphi^- = [u_2 - u_1 - 2\sqrt{K\varepsilon}]^+.$$

Note that $\varphi^\pm \in W_0^{1,2}(\Omega_j) \cap C(\overline{\Omega_j})$ and use $\psi_j^\pm = \varphi^\pm u_j$ as test functions in the weak formulation of $\Delta u_i + q_i u_i = 0$. We arrive at

$$\int_{\Omega_j} (|u_1| + |u_2|)(|u_1| - |u_2|)^2 \leq C \int_{\Omega_j} (|u_1| + |u_2|)\varepsilon,$$

Adding up w.r.t. j , and using the energy bound,

$$\int_{\Omega} ||u_1| - |u_2||^3 \leq \int_{\Omega} (|u_1| + |u_2|)(|u_1| - |u_2|)^2 \leq C\varepsilon,$$

Stability for $|u|$

W.l.o.g. we may assume $u_1, u_2 > 0$ in Ω_j . Set

$$\varphi^+ = [u_1 - u_2 - 2\sqrt{K\varepsilon}]^+, \varphi^- = [u_2 - u_1 - 2\sqrt{K\varepsilon}]^+.$$

Note that $\varphi^\pm \in W_0^{1,2}(\Omega_j) \cap C(\overline{\Omega_j})$ and use $\psi_j^\pm = \varphi^\pm u_j$ as test functions in the weak formulation of $\Delta u_j + q_j u_j = 0$. We arrive at

$$\int_{\Omega_j} (|u_1| + |u_2|)(|u_1| - |u_2|)^2 \leq C \int_{\Omega_j} (|u_1| + |u_2|)\varepsilon,$$

Adding up w.r.t. j , and using the energy bound,

$$\int_{\Omega} \||u_1| - |u_2|\|^3 \leq \int_{\Omega} (|u_1| + |u_2|)(|u_1| - |u_2|)^2 \leq C\varepsilon,$$

Integrability of $|u|^{-\delta}$

Lipschitz propagation of smallness (A. and Rosset '98, A., Rondi, Rosset and Vessella 2009) If

$$\frac{\int_{\Omega} |\nabla u|^2 + u^2}{\int_{\Omega} u^2} \leq \mathcal{F}$$

then for any $B_{\rho}(x_0) \subset \Omega$ we have

$$\int_{B_{\rho}(x_0)} u^2 \geq C \int_{\Omega} |\nabla u|^2 + u^2$$

where $C > 0$ only depends on ρ , K , Ω and on \mathcal{F} .

Note: Under our a priori assumptions:

$$\mathcal{F} = \frac{KE^2}{H^2}.$$

Integrability of $|u|^{-\delta}$

Lipschitz propagation of smallness (A. and Rosset '98, A., Rondi, Rosset and Vessella 2009) If

$$\frac{\int_{\Omega} |\nabla u|^2 + u^2}{\int_{\Omega} u^2} \leq \mathcal{F}$$

then for any $B_{\rho}(x_0) \subset \Omega$ we have

$$\int_{B_{\rho}(x_0)} u^2 \geq C \int_{\Omega} |\nabla u|^2 + u^2$$

where $C > 0$ only depends on ρ, K, Ω and on \mathcal{F} .

Note: Under our a priori assumptions:

$$\mathcal{F} = \frac{KE^2}{H^2}.$$

Integrability of $|u|^{-\delta}$

Doubling inequality (Garofalo and Lin '86) There exists $R = R(K)$ such that if $r_0 \leq R$ and $B_{r_0}(x_0) \subset \Omega$ then

$$\int_{B_{2r}(x_0)} u^2 \leq C \int_{B_r(x_0)} u^2, \forall r < \frac{r_0}{4}$$

where $C > 0$ only depends on r_0 and on the a priori data.

Integrability of $|u|^{-\delta}$

A_p property (Garofalo and Lin '86)

For any $d > 0$ there exist $p > 1$, $C > 0$, only depending on d and on the a priori data, such that for every $x_0 \in \Omega_d$ and every $r \leq d/4$

$$\left(\frac{1}{|B_r|} \int_{B_r(x_0)} u^2 \right) \left(\frac{1}{|B_r|} \int_{B_r(x_0)} u^{-\frac{2}{p-1}} \right)^{p-1} \leq C.$$

Concluding remarks

Quantitative estimates of unique continuation seem to be a necessary ingredient for stability estimates for IP with interior data, when available data depend on few solutions of the direct problem, and few restrictions can be imposed on such solutions.

Open issues may arise in investigating the vanishing rate of gradients or Jacobians.

Concluding remarks

Quantitative estimates of unique continuation seem to be a necessary ingredient for stability estimates for IP with interior data, when available data depend on few solutions of the direct problem, and few restrictions can be imposed on such solutions.

Open issues may arise in investigating the vanishing rate of gradients or Jacobians.

Concluding remarks

For equations with no zero order term, such as

$$\operatorname{div}(\sigma \nabla u) = 0 ,$$

with $\lambda^{-1}I \leq \sigma \leq \lambda I$, and if $n \geq 3$, $\sigma \in C^{0,1}$, estimates on the vanishing rate of $|\nabla u|^2$ are available.

If also a zero order term is present, e.g.:

$$\operatorname{div}(\sigma \nabla u) + qu = 0 ,$$

with $|q| \leq K$, estimates on the vanishing rate are known for $|\nabla u|^2 + |u|^2$.

The situation is **not** clear for $|\nabla u|^2$ alone.

Concluding remarks

For equations with no zero order term, such as

$$\operatorname{div}(\sigma \nabla u) = 0 ,$$

with $\lambda^{-1}I \leq \sigma \leq \lambda I$, and if $n \geq 3$, $\sigma \in C^{0,1}$, estimates on the vanishing rate of $|\nabla u|^2$ are available.

If also a zero order term is present, e.g.:

$$\operatorname{div}(\sigma \nabla u) + qu = 0 ,$$

with $|q| \leq K$, estimates on the vanishing rate are known for $|\nabla u|^2 + |u|^2$.

The situation is **not** clear for $|\nabla u|^2$ alone.

Concluding remarks

For equations with no zero order term, such as

$$\operatorname{div}(\sigma \nabla u) = 0 ,$$

with $\lambda^{-1}I \leq \sigma \leq \lambda I$, and if $n \geq 3$, $\sigma \in C^{0,1}$, estimates on the vanishing rate of $|\nabla u|^2$ are available.

If also a zero order term is present, e.g.:

$$\operatorname{div}(\sigma \nabla u) + qu = 0 ,$$

with $|q| \leq K$, estimates on the vanishing rate are known for $|\nabla u|^2 + |u|^2$.

The situation is **not** clear for $|\nabla u|^2$ alone.

Concluding remarks

An example

$$u_{xx} + qu = 0 \text{ on } (0, +\infty),$$

with

$$q(x) = \chi_{(0, \frac{\pi}{2})}(x),$$

and

$$u(x) = \begin{cases} \sin x & \text{if } 0 < x \leq \frac{\pi}{2}, \\ 1 & \text{if } x \geq \frac{\pi}{2}. \end{cases}$$

Concluding remarks

Further problems arise with Jacobians.

Example (Laugesen '96). $\forall \varepsilon > 0 \exists \Phi : \partial B \mapsto \partial B$
homeomorphism, such that $|\Phi(x) - x| < \varepsilon, \forall x \in \partial B$ and
the solution $U = (u_1, u_2, u_3)$ to

$$\begin{cases} \Delta U = 0, & \text{in } B, \\ U = \Phi, & \text{on } \partial B. \end{cases}$$

is **not** one-to-one.

Concluding remarks

Further problems arise with Jacobians.

Example (Laugesen '96). $\forall \varepsilon > 0 \exists \Phi : \partial B \mapsto \partial B$
homeomorphism, such that $|\Phi(x) - x| < \varepsilon, \forall x \in \partial B$ and
the solution $U = (u_1, u_2, u_3)$ to

$$\begin{cases} \Delta U = 0, & \text{in } B, \\ U = \Phi, & \text{on } \partial B. \end{cases}$$

is **not** one-to-one.

Concluding remarks

Example (Jin and Kazdan '91).

$\exists \sigma \in C^\infty(\mathbb{R}^3)$, $\lambda^{-1}I \leq \sigma \leq \lambda I$ and a solution $U = (u_1, u_2, u_3)$
to

$$\operatorname{div}(\sigma \nabla U) = 0 \text{ in } \mathbb{R}^3,$$

such that

$$\begin{cases} \det DU = 0, & \text{for } x_3 \leq 0, \\ \det DU > 0, & \text{for } x_3 > 0. \end{cases}$$

Introduction

An example

A priori
assumptions

Main Theorem

Stability for $|u|$

Quantitative
UCP

Concluding
remarks

End

The end.

THANKS!